

# **College Preparatory Integrated Mathematics Course II Notebook**

**Revised 09/23/2019**

**College Preparatory Integrated Mathematics Course II**  
**Learning Objective 1.1**  
**Section 5.2**

**Learning Objective 1.1: Define polynomial, monomial, binomial, trinomial, and degree. (Section 5.2 Objective 1)**

**Read Section 5.2 on page 322 and 323 in the textbook and answer the questions below.**

**Definitions**

1. A number or the product of a number and variables raised to powers is called \_\_\_\_\_.
2. The \_\_\_\_\_ of a term is the numerical factor of each term.
3. A \_\_\_\_\_ is a finite sum of terms of the form  $ax^n$ , where  $a$  is a real number and  $n$  is a whole number.
4. A \_\_\_\_\_ is a polynomial with exactly one term.
5. A \_\_\_\_\_ is a polynomial with exactly two terms.
6. A \_\_\_\_\_ is a polynomial with exactly three terms.
7. The \_\_\_\_\_ of a polynomial is the greatest \_\_\_\_\_ of any term of the polynomial.

**Example 1:** Find the degree of each term.

a)  $5y^3$

b)  $10xy$

c)  $z$

d)  $-3a^2b^5c$

e)  $8$

**Example 2:** Find the degree of each polynomial and tell whether the polynomial is a monomial, binomial, trinomial, or none of these.

a)  $5b^2 - 3b + 7$

b)  $7t + 3$

c)  $5x^2 + 3x - 6x^3 + 4$

d)  $1 - x^3 + x^4 + x$

**Learning Objective 1.1: Define polynomial functions (Section 5.2 Objective 2)**

**Read Section 5.2 on page 324 in the textbook and answer the questions below.**

**Example 3:** If  $P(x) = 2x^2 - 6x + 1$ , find the following.

a)  $P(1) =$

b)  $P(-3) =$

c)  $P(0) =$

**Learning Objective 1.1: Simplifying polynomials by combining Like Terms (Section 5.2 Objective 3)**

Read Section 5.2 on page 325 in the textbook and answer the questions below.

**Definitions**

1. Terms that contain exactly the same variables raised to exactly the same power called \_\_\_\_\_.

**Example 4:** Simplify each polynomial by combining any like terms.

a)  $-4y + 2y$

b)  $z + 5z^3$

c)  $15x^3 - x^3$

d)  $7a^2 - 5 - 3a^2 - 7$

e)  $\frac{3}{8}x^3 - x^2 + \frac{5}{6}x^4 + \frac{1}{12}x^3 - \frac{1}{2}x^4$

**Learning Objective 1.1: Add and Subtract polynomials (Section 5.2 Objective 4)**

Read Section 5.2 on page 327 in the textbook and answer the questions below.

**Definitions**

1. To add polynomials, combine all \_\_\_\_\_.
2. To subtract two polynomials, \_\_\_\_\_ the signs of the terms of the polynomial being subtracted and then add.

**Example 5:** Add or subtract.

a)  $(2x^2 + 7x + 6) + (x^2 - 6x^2 - 14)$

b)  $(-14x^3 - x + 2) + (-x^3 + 3x^2 + 4x)$

c)  $(8x^2 - 6x - 7) - (3x^2 - 5x)$

d)  $(2x - 5) - (7x^2 - 2x + 1)$

**College Preparatory Integrated Mathematics Course II**  
**Learning Objective 1.1**  
**Section 5.3**

**Learning Objective 1.1: Multiply monomials (Section 5.3 Objective 1)**

Read Section 5.3 on page 334 in the textbook and answer the questions below.

**Definitions**

1. To multiply exponential expressions with a common base, \_\_\_\_\_ exponents.

**Example 1:** Multiply.

a)  $5y \cdot 2y$

b)  $(5z^3) \cdot (-0.4z^5)$

c)  $(-\frac{1}{9}b^6) \cdot (-\frac{7}{8}b^3)$

**Learning Objective 1.1: Use the distributive property to multiply polynomials (Section 5.3 Objective 2)**

Read Section 5.3 on page 335 in the textbook and answer the questions below.

**Definitions**

1. To multiply two polynomials, multiply each term of the first polynomial by each term of the second polynomial and then combine \_\_\_\_\_.

**Example 2:** Multiply.

a)  $(x - 3)(x^2 - 6x + 1)$

b)  $(4a + 3b)^2$

c)  $(s + 2t)^3$

**Learning Objective 1.1: Multiply polynomials vertically (Section 5.3 Objective 3)**

Read Section 5.3 on page 337 in the textbook and answer the questions below.

**Example 3:** Find the product using a vertical format.

a)  $(5x^2 + 2x - 2)(x^2 - x + 3)$

c)  $(2 - x^2)(2x^2 + 4x - 1)$

**College Preparatory Integrated Mathematics Course II**  
**Learning Objective 1.1**  
**Section 5.4**

**Learning Objective 1.1: Multiply two binomial using the FOIL method. (Section 5.4 Objective 1)**  
 Read Section 5.4 on page 341 in the textbook an answer the questions below.

**Definitions**

**The FOIL method:**

1. F stands for the product of the \_\_\_\_\_ terms.
2. O stands for the product of the \_\_\_\_\_ terms.
3. I stands for the product of the \_\_\_\_\_ terms.
4. L stands for the product of the \_\_\_\_\_ terms.

**Example 1: Multiply.**

a)  $3(4x + 1)(5 - 2x)$

b)  $(4x - 1)^2$

**Learning Objective 1.1: Square a binomial (Section 5.4 Objective 2)**

Read Section 5.4 on page 342 in the textbook an answer the questions below.

**Definitions**

1.  $(a + b)^2 = a^2 + \underline{\hspace{2cm}} + b^2$
2.  $(a - b)^2 = a^2 - \underline{\hspace{2cm}} + b^2$

**Example 2: Use a special product to square each binomial.**

a)  $(b + 3)^2$

b)  $(x - y)^2$

d)  $(3y + 2)^2$

e)  $(a^2 - 5b)^2$

**Learning Objective 1.1: Multiplying the sum and difference of two terms. (Section 5.4 Objective 3)**

Read Section 5.4 on page 343 in the textbook and answer the questions below.

**Definitions**

1.  $(a + b)(a - b) = \underline{\hspace{2cm}}$ .

**Example 3:** Use a special product to multiply.

a)  $3(x + 5)(x - 5)$

b)  $(4b - 3)(4b + 3)$

c)  $(x + \frac{2}{3})(x - \frac{2}{3})$

d)  $(5s - t)(5s + t)$

e)  $(2y - 3z^2)(2y + 3z^2)$

**Learning Objective 1.1: Using special products (Section 5.4 Objective 4)**

Read Section 5.4 on page 344 in the textbook and answer the questions below.

**Example 4:** Use a special product to multiply, if possible. .

a)  $(4x + 3)(x - 6)$

b)  $(7b - 2)^2$

c)  $(x + 0.4)(x - 0.4)$

d)  $(x + 1)(x^2 + 5x - 2)$

e)  $(x^2 - \frac{3}{7})(3x^4 + \frac{2}{7})$

**College Preparatory Integrated Mathematics Course II**  
**Learning Objective 1.1**  
**Section 5.6**

**Learning Objective 1.1: Divide a polynomial by a monomial (Section 5.6 Objective 1)**  
 Read Section 5.6 on page 357 in the textbook and answer the questions below.

**Definitions**

1. Fractions that have a common denominator are added by adding the \_\_\_\_\_.

**Example 1:** Divide.

$$\frac{15x^4y^4 - 10xy + y}{5xy}$$

**Example 2:** In which of the following is  $\frac{x+5}{5}$  simplified correctly?

a)  $\frac{x}{5} + 1$

b)  $x$

c)  $x + 1$

**Learning Objective 1.1: Use long division to divide a polynomial by another polynomial (Section 5.6 Objective 2)**

Read Section 5.6 on page 358 in the textbook and answer the questions below.

**Definitions**

1. In  $18 \div 6 = 3$ , the 18 is the \_\_\_\_\_, the 3 is the \_\_\_\_\_, and the 6 is the \_\_\_\_\_.

**Example 3:** Divide.

a)  $x^3 + 27$  by  $x + 3$

b)  $x^2 + 2x - 6$  by  $x - 2$

**College Preparatory Integrated Mathematics Course II**  
**Learning Objective 1.1**  
**Section 5.7**

**Learning Objective 1.1: Use Synthetic division to divide a polynomial by a binomial (Section 5.7 Objective 1)**

Read Section 5.7 on page 364 in the textbook and answer the questions below.

**Definitions**

1. Which division problems are candidates for the synthetic division process?

a)  $(3x^2 + 5) \div (x + 4)$

c)  $(y^4 + y - 3) \div (x^2 + 1)$

b)  $(x^3 - x^2 + 2) \div (3x^3 - 2)$

d)  $x^5 \div (x - 5)$

**Example 1:** If  $P(x) = x^3 - 5x - 2$ ,

a) Find  $P(2)$  by substitution.

b) Use synthetic division to find the remainder when  $P(x)$  is divided by  $x - 2$ .

**Learning Objective 1.1: Using the Remainder Theorem (Section 5.7 Objective 2)**

Read Section 5.7 on page 366 in the textbook and answer the questions below.

**Definitions**

1. By Remainder Theorem, if a polynomial  $P(x)$  is divided by  $x - c$ , then the remainder is \_\_\_\_.

**Example 2:** Use the remainder theorem and synthetic division to find  $P(3)$  if

$$P(x) = 2x^5 - 18x^4 + 90x^2 + 59x$$



**College Preparatory Integrated Mathematics Course II**  
**Learning Objective 1.9**  
**Section 6.6**

**Learning Objective 1.9: Solve quadratic equations by factoring (Section 6.6 Objective 1)**  
Read Section 6.6 on page 417-422 in the textbook and answer the questions below.

**Definitions**

1. An equation that can be written in the form  $ax^2 + bx + c = 0$ , with  $a \neq 0$ , is called a \_\_\_\_\_ equation.
2. The form  $ax^2 + bx + c = 0$  is called the \_\_\_\_\_ of a quadratic equation.
3. If the product of two numbers is zero, then at least one of the numbers must be \_\_\_\_\_.
4. If  $a$  and  $b$  are real numbers and if  $a \cdot b = 0$ , then \_\_\_\_\_.

**Example 1:** Solve:

a)  $(x + 4)(x - 5) = 0$

b)  $(x - 12)(4x + 3) = 0$

c)  $x(7x - 6) = 0$

d)  $x^2 - 8x - 48 = 0$

e)  $9x^2 - 24x = -16$

f)  $x(3x + 7) = 0$

g)  $-3x^2 - 6x + 72 = 0$

**Learning Objective 1.9:** Solve equations with degree greater than 2 by factoring (Section 6.6 Objective 2)

Read Section 6.6 on page 421 in the textbook and answer the questions below.

**Example 2:** Solve:

a)  $7x^3 - 63x = 0$

b)  $(3x - 2)(2x^2 - 13x + 15) = 0$

c)  $5x^3 + 5x^2 - 30x = 0$

**Learning Objective 1.9:** Find x-intercepts of the graph of a quadratic equation in two variables. (Section 6.6 Objective 3)

Read Section 6.6 on page 423 in the textbook and answer the questions below.

**Definitions**

1. The graph of a quadratic equation in the form  $y = ax^2 + bx + c$  where  $a \neq 0$ , is called \_\_\_\_\_.

**Example 3:** Find the x-intercepts of the graph of  $y = x^2 - 6x + 8$ .

**Example 4:** Find the x-intercepts of the graph of  $y = x^2 + 4x + 4$ .

**Example 5:** Find the x-intercepts of the graph of  $y = 2x^2 + 2$ .

**College Preparatory Integrated Mathematics Course II**  
**Learning Objective 1.9**  
**Section 6.7**

**Learning Objective 1.9:** Solve problems that can be modeled by quadratic equations.(Section 6.7 Objective 1)

Read Section 6.7 on page 426-432 in the textbook and answer the questions below.

**Definitions**

1. In a right triangle, the side opposite the right angle is called the \_\_\_\_\_ .
2. In a right triangle, each side adjacent to the right angle is called a \_\_\_\_\_.
3. The Pythagorean theorem states that  $(leg)^2 + (leg)^2 = (\text{_____})^2$

**Example 1:** The square of a number minus eight times the number is equal to forty-eight. Find the number.

**Example 2:** Find two consecutive integers whose product is 41 more than their sum.

**Example 3:** Find the dimensions of a right triangle where the second leg is 1 unit less than double the first leg, and the hypotenuse is 1 unit more than double the length of the first leg.

**College Preparatory Integrated Mathematics Course II**  
**Learning Objective 1.3**  
**Section 7.2, 7.3, 7.4**

**Definitions**

1. If  $\frac{A}{B}$  and  $\frac{C}{D}$  are rational expressions, then  $\frac{A}{B} \cdot \frac{C}{D} =$  \_\_\_\_\_.
2. To divide two Rational Expressions, multiply the first rational expression by the \_\_\_\_\_ of the second rational expression.

**Example 1:** Multiply  $\frac{6x^2}{8x^3} \cdot \frac{16x}{12}$

**Example 2:** Multiply and simplify.  $\frac{(x-y)^2}{x+y} \cdot \frac{x}{x^2-xy}$

- a.) Re-write above Rational expression by Factoring all numerators and denominators
- b.) Multiply numerators and multiply denominators without distributing
- c.) Simplify by dividing out common factors.

**Example 3:** Divide  $\frac{4x^3y^7}{60} \div \frac{6x}{y^3}$

**Example 4:** Divide and simplify.  $\frac{10}{x^2-4} \div \frac{5x}{2x+4}$

- a.) Re-write above Rational expression by multiplying by Reciprocal of second rational expression
- b.) Factor all numerators and denominators and multiply remaining factors
- c.) Simplify by dividing out common factors.

**Example 5:** Divide.  $\frac{(x+3)^2}{4} \div \frac{4x+12}{16}$

### Learning Objective 1.3: Adding and Subtracting Rational Expressions with Common Denominators and Least Common Denominators

Read Section 7.3 on page 464 and answer the questions below.

#### Definitions

1. If  $\frac{A}{B}$  and  $\frac{C}{B}$  are rational expressions, then  $\frac{A}{B} + \frac{C}{B} = \frac{\quad}{B}$
2. If  $\frac{A}{B}$  and  $\frac{C}{B}$  are rational expressions, then  $\frac{A}{B} - \frac{C}{B} = \frac{\quad}{B}$
3. To add or subtract rational expressions, add or subtract \_\_\_\_\_ and place the sum or difference over the common denominator.
4. Use the distributive property to subtract  $2x - (x + 3) = \underline{\hspace{2cm}}$

**Example 6:** Add.  $\frac{5x-1}{4x} + \frac{2x-3}{4x}$

**Example 7:** Add.  $\frac{4m-3}{2m+7} + \frac{3m+8}{2m+7}$

**Example 8:** Subtract.  $\frac{8y}{y-3} - \frac{24}{y-3}$

**Example 9:** Subtract.  $\frac{3x}{x^2+3x-10} - \frac{6}{x^2+3x-10}$

**Example 10:** Subtract.  $\frac{7x+8}{9x+15} - \frac{5x-2}{9x+15}$

**Learning Objective 1.3: Adding and Subtracting Rational Expressions with Unlike Denominators**  
**Read Section 7.4 on page 472 and answer the questions below.**

**Definitions**

1. The least common denominator (LCD) is the product of all unique factors

2. If  $\frac{A}{B}$  and  $\frac{C}{D}$  are rational expressions, then  $\frac{A}{B} + \frac{C}{D} = \frac{\quad}{B}$

3. If  $\frac{A}{B}$  and  $\frac{C}{D}$  are rational expressions, then  $\frac{A}{B} - \frac{C}{D} = \frac{\quad}{B}$

**Four Steps to Adding and Subtracting Rational Expressions with Unlike Denominators.**

Step 1: Find the LCD of all the rational expressions.

Step 2: Rewrite each rational expression as an equivalent expression whose denominator is the LCD found in Step 1.

Step 3: Add or subtract numerators and write the sum or difference over the common

denominator.

Step 4: Simplify or write the rational expression in simplest form.

**Example 11:** Add.  $\frac{15}{7a} + \frac{8}{6a} =$

**Example 12:** Add.  $4 + \frac{4}{x}$

**Example 13:** Add.  $\frac{4}{x^2 - x - 6} + \frac{x}{x^2 + 5x + 6}$

**Example 14:** Add.  $\frac{9}{x^2 + 5x - 6} + \frac{6}{x + 6}$

**Example 15:** Subtract.  $\frac{7}{2x - 3} - 3$

**Example 16:** Subtract.  $1 - \frac{1}{x}$

**Example 17:** Subtract.  $\frac{5}{2x-6} - \frac{3}{6-2x}$

**Example 18:** Subtract.  $\frac{x^2}{x} - \frac{2x+8}{2x}$



**College Preparatory Integrated Mathematics Course I**  
**Learning Objective 1.4**  
**Section 7.7**

**Learning Objective 1.4: Simplifying Complex Fractions**

Read Section 7.7 on page 499 and answer the questions below.

**Definitions**

**Method 1: Simplifying a Complex Fraction**

**Step 1:** Simplify the numerator and the denominator of the complex fraction so that each is a single fraction.

**Step 2:** Perform the indicated division by multiplying the numerator of the complex fraction by the \_\_\_\_\_ of the denominator of the complex fraction.

**Step 3:** Simplify if possible

**Method 2: Simplifying a Complex Fraction**

**Step 1:** Multiply the numerator and the denominator of the complex fraction by the \_\_\_\_\_ of the fractions in both the numerator and the denominator.

**Step 2:** Simplify

**Example 1:** Use Method 1 above to simplify.

$$\frac{1 + \frac{1}{x}}{4 - \frac{4}{x}}$$

Step1:

Step2:

Step3:

**Example 2:** Use Method 1 above to simplify.  $\frac{\frac{x}{2} + 2}{\frac{x}{4} - 4}$

**Example 3:** Use Method 2 above to simplify.  $\frac{\frac{6x^2}{8x^3}}{\frac{12}{16x}}$

Step1:

Step2:

**Example 4:** Use Method 2 above to simplify.  $\frac{\frac{1}{y^2} + \frac{2}{3}}{\frac{1}{y} - \frac{5}{6}}$

**College Preparatory Integrated Mathematics Course I**  
**Learning Objective 1.6**  
**Section 10.2 and 10.3**

**Learning Objective 1.6: Simplifying Rational Exponents**

Read Section 10.2 on page 605 and answer the questions below.

**Definitions**

1. If  $n$  is a positive integer greater than 1, then fill in the blank  $a^{\frac{1}{n}} = \sqrt[n]{a}$
2. If  $m$  and  $n$  are positive integers greater than 1, with  $m/n$  in simplest form, then fill in the blanks :  $a^{\frac{m}{n}} = \sqrt[n]{a^m} = \left(\sqrt[n]{a}\right)^m$
3. If  $a^{\frac{m}{n}}$  is a nonzero real number, then fill in the blanks:  $a^{-\frac{m}{n}} = \frac{1}{a^{\frac{m}{n}}}$

**Example 1:** Use radical notation to write the following. Simplify if possible.  $81^{\frac{1}{4}}$

**Example 2:** Use radical notation to write the following. Simplify if possible.  $(32x^{10})^{1/5} =$

**Example 3:** Use radical notation to write the following. Simplify if possible.  $-(16x^8)^{\frac{1}{2}}$

**Learning Objective 1.6: Simplifying Radical Expressions**

Read Section 10.3 on page 612 and answer the questions below.

**Definitions**

1. Product Rule for Radicals: Fill in the blank  $\sqrt[n]{a} \cdot \sqrt[n]{b} = \sqrt[n]{\quad}$

2. Quotient Rule for Radicals: Fill in the blanks:  $\sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{\quad}}{\sqrt[n]{\quad}}$

**Example 4:** Use rational exponents to write as a single radical.  $\sqrt[3]{5} \cdot \sqrt{2} =$

**Example 5:** Use rational exponents to write as a single radical and Simplify.  $\sqrt[3]{-343x^6}$

**Example 6:** Multiply and Simplify.  $\sqrt{\frac{2}{a}} \cdot \sqrt{\frac{b}{3}}$

**Example 7:** Simplify.  $\sqrt[3]{\frac{8}{27}}$

**Example 8:** Use the quotient rule to divide, and simplify if possible  $\sqrt{\frac{20}{x^{16}}} =$

**College Preparatory Integrated Mathematics Course II**  
**Learning Objective 1.7**  
**Section 10.4**

**Learning Objective 1.7: Add or subtract radical expressions (Section 10.4 Objective 1)**

Read Section 10.4 on page 620 in the textbook and answer the questions below.

**Definitions**

1. Radicals with the same index and the same radicand are \_\_\_\_\_.

**Example 1:** Add or subtract. Assume that variables represent positive real numbers.

a)  $3\sqrt{17} + 5\sqrt{17}$

b)  $7\sqrt[3]{5z} - 12\sqrt[3]{5z}$

**Example 2:** Add or subtract. Assume that variables represent positive real numbers.

a)  $\sqrt{24} + 3\sqrt{54}$

b)  $\sqrt[3]{24} - 4\sqrt[3]{81} + \sqrt[3]{3}$

c)  $\sqrt{75x} - 3\sqrt{27x} + \sqrt{12x}$

d)  $\frac{\sqrt{28}}{3} - \frac{\sqrt{7}}{4}$

**Learning Objective 1.7: Multiply radical expressions (Section 10.4 Objective 2)**

Read Section 10.4 on page 623 in the textbook and answer the questions below.

**Example 3:** Multiply.

a)  $\sqrt{5}(2 + \sqrt{15})$

b)  $(\sqrt{2} - \sqrt{5})(\sqrt{6} + 2)$

c)  $(\sqrt{6} - 3)^2$

d)  $(3\sqrt{z} - 4)(2\sqrt{z} + 3)$

$$e)(\sqrt{x+2} + 3)^2$$

**College Preparatory Integrated Mathematics Course II**  
**Learning Objective 1.7**  
**Section 10.5**

**Learning Objective 1.7: Rationalize denominators (Section 10.5 Objective 1)**

Read Section 10.5 on page 626 in the textbook and answer the questions below.

**Definitions**

1. The process of writing an equivalent expression, but without a radical in the denominator is called \_\_\_\_\_.

**Example 1:** Rationalize the denominator of each expression.

a)  $\frac{5}{\sqrt{3}}$

b)  $\frac{3\sqrt{25}}{\sqrt{4x}}$

c)  $\sqrt[3]{\frac{2}{9}}$

**Learning Objective 1.7: Rationalize denominators having two terms (Section 10.5 Objective 2)**

Read Section 10.5 on page 619 in the textbook and answer the questions below.

**Definitions**

1. Two expressions  $a + b$  and  $a - b$  are called \_\_\_\_\_.

**Example 2:** Rationalize the denominator.

a)  $\frac{5}{3\sqrt{5}+2}$

b)  $\frac{\sqrt{2}+5}{\sqrt{3}-\sqrt{5}}$

c)  $\frac{3\sqrt{x}}{2\sqrt{x}+\sqrt{y}}$

**Learning Objective 1.7: Rationalize numerators (Section 10.5 Objective 3)**

Read Section 10.5 on page 629 in the textbook and answer the questions below.

**Definitions**

1. The process of writing an equivalent expression, but without a radical in the numerator is called \_\_\_\_\_.

**Example 3:** Rationalize numerator.

a)  $\frac{\sqrt{32}}{\sqrt{80}}$

b)  $\frac{\sqrt[3]{5b}}{\sqrt[3]{2a}}$

c)  $\frac{\sqrt{x}-3}{4}$

**College Preparatory Integrated Mathematics Course I**  
**Learning Objective 1.7**  
**Section 10.6**

**Learning Objective 1.7: Simplifying Radical Expressions and Solve Radical Equations**

Read Section 10.6 on page 633 and answer the questions below.

**Definitions**

1. **Power Rule:** Fill in the blanks: If both sides of an equation are raised to the same power, \_\_\_\_\_ solutions of the original equation are *among* the solutions of the \_\_\_\_\_ equation.
2. **Pythagorean Theorem:** If  $a$  and  $b$  are lengths of the legs of a right triangle and  $c$  is the length of the hypotenuse, then fill in the blanks: \_\_\_\_\_ + \_\_\_\_\_ = \_\_\_\_\_

**Example 1:** Solve.  $\sqrt{x+1} = 5$

**Example 2:** Solve  $x\sqrt{2} = \sqrt{9}$

**Example 3:** Solve.  $2x + \sqrt{x+1} = 8$

**Example 4:** Solve.  $\sqrt{5x} = -5$

**Example 5:** Solve.  $\sqrt{y+5} = 2 - \sqrt{y-4}$

**Example 6:** Find the length of the hypotenuse of a right triangle when the length of the two legs are 2 inches and 7 inches.

**Example 7:** Find the length of the leg of a right triangle. Give the exact length and a two-decimal-approximation. Let  $a = 2$  meters and  $c = 9$  meters



**College Preparatory Integrated Mathematics Course II**  
**Learning Objective 1.9**  
**Section 11.1**

**Learning Objective 1.9: Use the square root property to solve quadratic equations.(Section 11.1 Objective 1)**

**Read Section 11.1 on page 661 in the textbook an answer the questions below.**

**Definitions**

1. A \_\_\_\_\_ equation is an equation that can written in the form  $x^2 + bx + c$ .
2. If  $b$  is a real number and if  $a^2 = b$ , then  $a =$ \_\_\_\_\_.

**Example 1:** Use square root property to solve equations.

**h)  $x^2 = 32$**

**b)  $5x^2 - 50 = 0$**

**c)  $(x + 3)^2 = 20$**

**d)  $(5x - 2)^2 + 2 = -7$**

**Learning Objective 1.9: Solving by completing the square (Section 11.1 Objective 2)**

**Read Section 11.1 on page 654 in the textbook an answer the questions below.**

**Definitions**

1. The process of writing a quadratic equation so that one side is a perfect square trinomial is called \_\_\_\_\_.
2. A perfect square trinomial is one that can be factored as a \_\_\_\_\_ squared.
3. To solve  $x^2 + 6x = 10$  by completing the square, add \_\_\_\_\_ to both sides.
4. To solve  $x^2 + bx = c$  by completing the square, add \_\_\_\_\_ to both sides.

**Example 2:** Solve equations by completing the square.

**a)  $b^2 + 4b = 3$**

b)  $2x^2 - 5x + 7 = 0$

c)  $3x^2 - 12x + 1 = 0$

**Learning Objective 1.9: Solving problems modeled by quadratic equations (Section 11.1 Objective 3)**

**Read Section 11.1 on page 666 in the textbook and answer the questions below.**

**Definitions**

2. The formula  $I = Prt$  is a formula for \_\_\_\_\_.
3. The interest computed on money borrowed or money deposited is \_\_\_\_\_.

**Example 3:** Use the formula  $A = P(1 + r)^t$  to find the interest rate  $r$  if \$5000 compounded annually grows to \$5618 in 2 years.

**College Preparatory Integrated Mathematics Course II**  
**Learning Objective 1.9**  
**Section 11.2**

**Learning Objective 1.9: Solve quadratic equations by using the quadratic formula.(Section 11.2 Objective 1)**

**Read Section 11.2 on page 671 in the textbook an answer the questions below.**

**Definitions**

1. The quadratic equation written in the form  $x^2 + bx + c = 0$  , when  $a \neq 0$  has the solutions \_\_\_\_\_.

**Example 1:** Solve equations by using quadratic formula.

a)  $3x^2 - 5x - 2 = 0$

b)  $3x^2 - 8x = 2$

c)  $\frac{1}{8}x^2 - \frac{1}{4}x - 2 = 0$

**Learning Objective 1.9: Determine the number and type of solutions of a quadratic equation by using the discriminant.(Section 11.2 Objective 2)**

**Read Section 11.2 on page 674 in the textbook an answer the questions below.**

**Definitions**

1. The radicand  $b^2 - 4ac$  is called \_\_\_\_\_.
2. If  $b^2 - 4ac$  is positive, the quadratic equation has \_\_\_\_\_ solutions.
3. If  $b^2 - 4ac$  is zero, the quadratic equation has \_\_\_\_\_ solutions.
4. If  $b^2 - 4ac$  is negative, the quadratic equation has \_\_\_\_\_ solutions.

**Example 2:** Use the discriminant to determine the number and type of solutions of each quadratic equation.

a)  $x^2 - 6x + 9 = 0$

b)  $x^2 - 3x - 1 = 0$

c)  $7x^2 + 11 = 0$

**Learning Objective 1.9: Solve problems modeled by quadratic equations.(Section 11.2 Objective 3)**

**Read Section 11.2 on page 675 in the textbook and answer the questions below.**

**Example 3:** A toy rocket is shot upward from the top of a building, 45 feet high, with an initial velocity of 20 feet per second. The height  $h$  in feet of the rocket after  $t$  seconds is

$$h = -16t^2 + 20t + 45$$

How long after the rocket is launched will it strike the ground? Round to the nearest tenth of a second.

**College Preparatory Integrated Mathematics Course II**  
**Learning Objective 1.9**  
**Section 11.3**

**Learning Objective 1.9: Solve various equations that are quadratic in form.(Section 11.3 Objective 1)**

**Read Section 11.3 on page 681 in the textbook and answer the questions below.**

**Definitions**

1. The best way to solve the quadratic equation in the form  $(ax + b)^2 = c$  is \_\_\_\_\_.

**Example 1:** Solve:

a)  $x - \sqrt{x + 1} - 5 = 0$

b)  $\frac{5x}{x+1} - \frac{x+4}{x} = \frac{3}{x(x+1)}$

c)  $p^4 - 7p^2 - 144 = 0$

d)  $(x - 3)^2 - 3(x - 3) - 4 = 0$

**Learning Objective 1.9: Solve problems that lead to quadratic equations.(Section 11.3 Objective 2)**

**Read Section 11.3 on page 684 in the textbook and answer the questions below.**

**Definitions**

1. Four steps to solve a word problem are \_\_\_\_\_, \_\_\_\_\_, \_\_\_\_\_, and \_\_\_\_\_.

**Example 2:** Together, Katy and Steve can groom all the dogs at the Barkin' Doggies Day Care in 4 hours. Alone, Katy can groom the dogs 1 hour faster than Steve can groom the dogs alone. Find the time in which each of them can groom the dogs alone.

**College Preparatory Integrated Mathematics Course II**  
**Learning Objective 2.1**  
**Section 11.4**

**Learning Objective 2.1: Solve polynomial inequalities of degree 2 or more.(Section 11.4 Objective 1)**

**Read Section 11.4 on page 691 in the textbook and answer the questions below.**

**Definitions**

1. A \_\_\_\_\_ is an inequality that can be written so that one side is a quadratic expression and the other side is 0.
2. An inequality is written in standard form if one side is an \_\_\_\_\_ and the other side is \_\_\_\_\_.

**Example 1: Solve inequalities.**

**a)  $(x - 4)(x + 3) > 0$**

**b)  $x^2 - 8x \leq 0$**

**c)  $(x + 3)(x - 2)(x + 1) \leq 0$**

**Learning Objective 2.1: Solve inequalities that contain rational expressions with variables in the denominator.(Section 11.4 Objective 2)**

**Read Section 11.4 on page 694 and 695 in the textbook and answer the questions below.**

**Definitions**

1. The first step to solve a rational inequality is solve for values that make all denominators \_\_\_\_\_.
2. An \_\_\_\_\_ **interval** does not include its endpoints, and is indicated with parentheses.
3. A \_\_\_\_\_ **interval** includes its endpoints, and is denoted with square brackets.

**Example 2: Solve inequalities.**

a)  $\frac{x-5}{x+4} \leq 0$

b)  $\frac{7}{x+3} < 5$





**College Preparatory Integrated Mathematics Course II**  
**Learning Objective 2.1**  
**Section 11.5**

**Learning Objective 2.1: Graph Quadratic Functions and Inequalities**

Read Section 11.5 on page 698 and answer the questions below.


**Definitions**

1. A \_\_\_\_\_ is a function that can be written in the form  $f(x) = ax^2 + bx + c$ , where  $a$ ,  $b$ , and  $c$  are real numbers and  $a \neq 0$ .
2. If  $a > 0$ , the parabola opens \_\_\_\_\_.
3. If  $a < 0$ , the parabola opens \_\_\_\_\_.
4. The \_\_\_\_\_ of a parabola is the \_\_\_\_\_ point if the graph opens upward and the \_\_\_\_\_ point if the parabola opens downward.
5. The \_\_\_\_\_ is the vertical line that passes through the vertex.



**Example 1:** Graph each quadratic function on the same coordinate plane. Label the vertex and sketch and label the axis of symmetry.

- a.  $f(x) = x^2$
- b.  $f(x) = x^2 + 2$
- c.  $f(x) = x^2 - 3$




**Definition:** Graphing the Parabola Defined by  $f(x) = x^2 + k$

1. If  $k$  is positive, the graph of  $f(x) = x^2 + k$  is the graph of  $y = x^2$  shifted \_\_\_\_\_.
2. If  $k$  is negative, the graph of  $f(x) = x^2 + k$  is the graph of  $y = x^2$  shifted \_\_\_\_\_.
3. The vertex is \_\_\_\_\_ and the axis of symmetry is \_\_\_\_\_.

**Example 2:** Graph each quadratic function on the same coordinate plane. Label the vertex and sketch and label the axis of symmetry.

- a.  $f(x) = x^2$
- b.  $f(x) = (x - 2)^2$
- c.  $f(x) = (x + 3)^2$



**Definition:** Graphing the Parabola Defined by  $f(x) = (x - h)^2$

1. If  $h$  is positive, the graph of  $f(x) = (x - h)^2$  is the graph of  $y = x^2$  shifted to the \_\_\_\_\_.

2. If  $h$  is negative, the graph of  $f(x) = (x - h)^2$  is the graph of  $y = x^2$  shifted to the \_\_\_\_\_.

3. The vertex is \_\_\_\_\_ and the axis of symmetry is \_\_\_\_\_.


Definition: Graphing the Parabola Defined by  $f(x) = (x - h)^2 + k$

1. The parabola has the same shape as \_\_\_\_\_.


2. The vertex is \_\_\_\_\_ and the axis of symmetry is \_\_\_\_\_.

**Example 3:** Graph each quadratic function. Label the vertex and sketch and label the axis of symmetry.

a.  $f(x) = (x - 2)^2 + 1$




b.  $f(x) = (x + 1)^2 - 3$



**Example 4:** Graph each quadratic function on the same coordinate plane. Label the vertex and sketch and label the axis of symmetry.

- $f(x) = x^2$
- $f(x) = 2x^2$
- $f(x) = \frac{1}{2}x^2$




**Definition:** Graphing the Parabola Defined by  $f(x) = ax^2$

- If  $|a| > 1$ , the graph of the parabola is \_\_\_\_\_ than the graph of  $y = x^2$ .
- If  $|a| < 1$ , the graph of the parabola is \_\_\_\_\_ than the graph of  $y = x^2$ .

**Example 5:** Graph each quadratic function on the same coordinate plane. Label the vertex and sketch and label the axis of symmetry.


- $f(x) = x^2$
- $f(x) = -x^2$



**Definition:** Graph of a Quadratic Function


- The graph of a quadratic function written in the form  $f(x) = a(x - h)^2 + k$  is a parabola with vertex \_\_\_\_\_.
- If  $a > 0$ , the parabola opens \_\_\_\_\_.

3. If  $a < 0$ , the parabola opens \_\_\_\_\_.
4. The axis of symmetry is the line whose equation is \_\_\_\_\_.




**Example 6:** Graph each quadratic function. Label the vertex and two other points on the graph. Sketch and label the axis of symmetry.

a.  $f(x) = -2(x - 3)^2 + 4$



b.  $f(x) = \frac{1}{3}(x + 3)^2 - 2$



**College Preparatory Integrated Mathematics Course II**  
**Learning Objective 2.1**  
**Section 11.6**

**Learning Objective 2.1: Graph Quadratic Functions and Inequalities**


Read Section 11.6 on page 706 and answer the questions below.

**Definitions**


1. The graph of a quadratic function is a \_\_\_\_\_.
2. To write a quadratic function in the form  $f(x) = a(x - h)^2 + k$ , we

\_\_\_\_\_.


**Example 1:** Graph  $f(x) = x^2 + 6x + 9$ . Find the vertex and any intercepts.



**Example 2:** Graph  $f(x) = -2x^2 + 4x + 6$ . Find the vertex and any intercepts.



**Example 3:** Graph  $f(x) = x^2 + x + 6$ . Find the vertex and any intercepts.




**Example 4:** Complete the square on  $y = ax^2 + bx + c$  and write the equation in the form  $y = a(x - h)^2 + k$

**Definition: Vertex Formula**

1. The graph of  $f(x) = ax^2 + bx + c$ , when  $a \neq 0$ , is a parabola with vertex \_\_\_\_\_.


**Example 5:** Find the vertex of the graph of each quadratic function. Determine whether the graph opens upward or downward, find any intercepts, and graph the function.

a.  $f(x) = x^2 + 5x + 4$





b.  $f(x) = x^2 - 4x + 4$



**Definition:** Minimum and Maximum Values

1. The quadratic function whose graph is a parabola that opens upward has a \_\_\_\_\_.
2. The quadratic function whose graph is a parabola that opens downward has a \_\_\_\_\_.
3. The \_\_\_\_\_ of the vertex is the minimum or maximum value of the function.

**Example 6:** An arrow is fired into the air with an initial velocity of 96 feet per second. The height in feet of the arrow  $t$  seconds after it was shot into the air is given by the function  $h(x) = -16t^2 + 96t$ . Find the maximum height of the arrow.

# College Preparatory Integrated Mathematics Course II

## Learning Objective 3.1

### Section 2.4

#### Learning Objective 3.1: Solve Word Problems

Read Section 2.4 on page 106 and answer the questions below.

**Definitions:** General Strategy for Problem Solving

1. UNDERSTAND the problem. Some ways of doing this are to:
  - 
  - 
  - 
  -
2. TRANSLATE the problem into an equation.
3. SOLVE the equation.
4. INTERPRET the result: *Check* the proposed solutions in the stated problem and state your conclusion.

**Example 1 – Solving Direct Translation Problems:** Eight is added to a number and the sum is doubled. The result is 11 less than the number. Find the number.

**Example 2 – Solving Direct Translation Problems:** Three times the difference of a number and 2 is equal to 8 subtracted from twice a number. Find the integers.

**Example 3 – Solving Problems Involving Relationships Among Unknown Quantities:** A 22-ft pipe is cut into two pieces. The shorter piece is 7 feet shorter than the longer piece. What is the length of the longer piece?

**Example 4 – Solving Problems Involving Relationships Among Unknown Quantities:** A college graduating class is made up of 450 students. There are 206 more girls than boys. How many boys are in the class?

**Example 5 – Solving Consecutive Integer Problems:** The room numbers of two adjacent hotel rooms are two consecutive odd numbers. If their sum is 1380, find the hotel room numbers.

# College Preparatory Integrated Mathematics Course II

## Learning Objective 3.1

### Section 2.5

#### Learning Objective 3.1: Solve Word Problems

Read Section 2.5 on page 117 and answer the questions below.

#### Definitions

1. An equation that describes a known relationship among quantities, such as distance, time, volume, weight, and money, is called a \_\_\_\_\_.
2. These quantities are represented by \_\_\_\_\_ and are thus \_\_\_\_\_ of the formula.

#### Common Formulas

<i>Formulas</i>	<i>Their Meanings</i>
$A = lw$	
$I = PRT$	
$P = a + b + c$	
$d = rt$	
$V = lwh$	
$F = \left(\frac{9}{5}\right)C + 32$ or $F = 1.8C + 32$	

**Example 1 – Using Formulas to Solve Problems:** Substitute the given values into each given formula and solve for the unknown variable. If necessary, round to one decimal place.

- a. Distance Formula  
 $d = rt$ ;  $t = 9$ ,  $d = 63$

- b. Perimeter of a rectangle  
 $P = 2l + 2w$ ;  $P = 32$ ,  $w = 7$

- c. Volume of a pyramid  
 $V = \frac{1}{3}Bh$ ;  $V = 40$ ,  $h = 8$

- d. Simple interest  
 $I = prt$ ;  $I = 23$ ,  $p = 230$ ,  $r = 0.02$

**Example 2 – Using Formulas to Solve Problems:** Convert the record high temperature of  $102^{\circ}\text{F}$  to Celsius.

**Example 3 – Using Formulas to Solve Problems:** You have decided to fence an area of your backyard for your dog. The length of the area is 1 meter less than twice the width. If the perimeter of the area is 70 meters, find the length and width of the rectangular area.

**Example 4 – Using Formulas to Solve Problems:** For the holidays, Christ and Alicia drove 476 miles. They left their house at 7 a.m. and arrived at their destination at 4 p.m. They stopped for 1 hour to rest and re-fuel. What was their average rate of speed?

**Example 5 – Solving a Formula for One of Its Variables:** Solve each formula for the specified variable.

a. Area of a triangle

$$A = \frac{1}{2}bh \text{ for } b$$

b. Perimeter of a triangle

$$P = s_1 + s_2 + s_3 \text{ for } s_3$$

c. Surface area of a special rectangular box

$$S = 4lw + 2wh \text{ for } l$$

d. Circumference of a circle

$$C = 2\pi r \text{ for } r$$

# College Preparatory Integrated Mathematics Course II

## Learning Objective 3.1

### Section 2.6

#### Learning Objective 3.1: Solve Word Problems

Read Section 2.6 on page 128 and answer the questions below.

Review: General Strategy for Problem Solving

1. UNDERSTAND the problem.
2. TRANSLATE the problem into an equation.
3. SOLVE the problem.
4. INTERPRET the results: *Check* the proposed solution in the stated problem and *state* your conclusion.

**Example 1 – Solving Percent Equations:** Find each number described.

- a. 5% of 300 is what number?
- b. 207 is 90% of what number?
- c. 15 is 1% of what number?
- d. What percent of 350 is 420?

**Example 2 – Solving Discount and Mark-up Problems:** A “Going-Out-Of-Business” sale advertised a 75% discount on all merchandise. Find the discount and the sale price of an item originally priced at \$130. If needed, round answers to the nearest cent.

**Example 3 – Solving Discount and Mark-up Problems:** Recently, an anniversary dinner cost \$145.23 excluding tax. Find the total cost if a 15% tip is added to the cost.

**Example 4 – Solving Percent Increase and Percent Decrease Problems:** In 2004, a college campus had 8,900 students enrolled. In 2005, the same college campus had 7,600 students enrolled. Find the percent decrease. Round to the nearest whole percent.

**Example 5 – Solving Mixture Problems:** How much pure acid should be mixed with 4 gallons of a 30% acid solution in order to get an 80% acid solution? Use the following table to model the situation.

	Number of Gallons · Acid Strength = Amount of Acid		
Pure Acid			
30% Acid Solution			
80% Acid Solution Needed			

# College Preparatory Integrated Mathematics Course II

## Learning Objective 4.1

### Section 8.2

**Learning Objective 3.1: Recognize functional notation and evaluate functions.**

**Read Section 8.2 on page 525 and answer the questions below.**

**Definition:** (Review from Section 3.6, pg. 229)

1. A \_\_\_\_\_ is a set of ordered pairs that assigns to each  $x$ -value exactly one  $y$ -value.
2. The variable  $x$  is the \_\_\_\_\_ because any value in the domain can be assigned to  $x$ .
3. The variable  $y$  is the \_\_\_\_\_ because its value depends on  $x$ .
4. The symbol  $f(x)$  means \_\_\_\_\_ and is read " $f$  of  $x$ ." This is called function notation and  $y = f(x)$ .

**Example 1:** For each given function value, write a corresponding ordered pair.

a.  $f(3) = 6$

b.  $g(0) = -\frac{1}{2}$

c.  $h(-2) = 9$

**Example 2:** Use the graph of the following function  $f(x)$  to find each value. Write the corresponding ordered pair for each.


a.  $f(1) =$

b.  $f(-3) =$

c.  $f(0) =$

d. Find  $x$  such that  $f(x) = 2$ .

e. Find  $x$  such that  $f(x) = 0$ .





**Example 2:** For each function, find the value of  $f(-3)$ ,  $f(2)$ , and  $f(0)$ . Then write the corresponding ordered pairs.

a.  $f(x) = -\frac{1}{3}x - 5$

$$f(-3) =$$

$$f(2) =$$

$$f(0) =$$

b.  $f(x) = 3x^2 - 2x - 2$

$$f(-3) =$$

$$f(2) =$$

$$f(0) =$$

c.  $f(x) = |-3 - x|$

$$f(-3) =$$

$$f(2) =$$

$$f(0) =$$